

Fact Sheet

Hydration and the Active Child



In hot weather it can be very easy for physically active children to suffer from dehydration, heat exhaustion, or even heat stroke. A physically active child can be described as a child who participates in at least an hour of moderate intensity physical activity every day. Some examples of moderate intensity activity include team sports such as soccer, football or netball, swimming training, running or tennis.

It is known that children do not stop to drink as much as adults do and are likely to only stop when they are quite thirsty. At this stage it is likely that children are already mildly dehydrated.

What is the best thing to drink?

- Water should always be the first choice of fluid. Children should be encouraged to keep a water bottle with them at all times and to stop regularly during activity to re-hydrate.
- Drinks that also contain some carbohydrate and extra salts such as sports drinks may have a small additional benefit over water for children engaged in high intensity activity of over an hour's duration.
- During hot weather, and during high intensity exercise, a good option would be for children to

have both chilled water and sports drink on hand to encourage rehydration.

- When choosing sports drinks, be aware that some "energy" type drinks may also contain other additives such as caffeine. These are not suitable for children.

How much to drink?

- It is somewhat difficult to accurately estimate how much fluid an active child requires on any one day. This is due to variations in the intensity of exercise and the weather.
- A child aged under 10 years will require about 200mls of fluid prior to starting activity and at least 100mls for every 20 mins they are involved in physical activity. This is equal to ~ a 300ml bottle of water or sports drink per hour.
- Children aged 10 years and above will need double this volume of fluid to keep adequately hydrated.

Remember

- Active children should be encouraged to drink regularly to make sure they do not feel thirsty. Regular drink breaks during activity are therefore vital.

This fact sheet is for education purposes only.
Please consult with your doctor or other health professional
to make sure this information is right for your child.

This document was reviewed on Tuesday, 19th December 2006.

the
children's
hospital at Westmead

www.chw.edu.au

 **SYDNEY
CHILDREN'S
HOSPITAL
RANDWICK**

www.sch.edu.au


Kaleidoscope
HUNTER CHILDREN'S HEALTH NETWORK

www.kaleidoscope.org.au